

Flow Cytometry (CMS Article A56464)

CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367

Medicare Local Coverage Determination (LCD)

The list of ICD10 codes provided consists of *commonly utilized diagnosis codes*.

- It is the responsibility of the ordering provider to ensure appropriate diagnostic coding for a test.
- If you are ordering this test for diagnostic reasons that are not covered under Medicare policy, an Advanced Beneficiary Notice (ABN) form is required.
- To view the CMS Local Coverage Determination for Flow Cytometry, visit the following website:
[Article - Billing and Coding: Flow Cytometry \(A56464\) \(cms.gov\)](#)

Coverage Indications, Limitations, and/or Medical Necessity

Coverage Indications, Limitations, and/or Medical Necessity

Flow cytometry (FCM) is a complex process to examine blood, body fluids, cerebrospinal fluid (CSF), bone marrow, lymph node, tonsil, spleen and other solid tissues. The use of peripheral blood and fine needle aspirate material avoids more invasive procedures for diagnosis.

A flow cytometer evaluates the physical and/or chemical characteristics of single cells as the cells pass individually in a fluid stream through a measuring device. Surface receptors, intracellular molecules, and DNA bind with fluorescent dyes that allow detection and evaluation.

When light of one wave length excites electrons of certain chemicals to energy levels above their ground state and upon return to ground state emits light of a longer wavelength, fluorescence is produced. A flow cytometer detects cell characteristics by measuring the fluorescence produced by fluorochromes conjugated either directly with cell components or conjugated to antibodies directed against cell components.

Indications

Cytopenias and Hypercellular Hematolymphoid Disorders

Hematolymphoid neoplasia can present with cytopenias (anemia, leukopenia and/or thrombocytopenia) or elevated leukocyte counts. If medical review and preliminary laboratory testing fails to reveal a cause, bone marrow aspiration and biopsy are indicated to rule out an infiltrative process or a stem cell disorder. FCM is essential to evaluate hematolymphoid lineages. Although anemia commonly occurs in nonneoplastic diseases, anemia alone should not automatically trigger FCM.

FCM may be useful in hypercellular hematolymphoid disorders to differentiate reactive conditions from neoplastic conditions. In the absence of blasts, neutrophilic leukocytosis is not generally an indication for FCM. Isolated

REMINDER: The ordering provider is solely responsible for assigning diagnosis (codes). PDL does not, through this Reference Guide or otherwise, recommend any particular diagnosis codes. PDL will submit to Medicare only the diagnosis (codes) provided to PDL by the ordering provider and/or his/her authorized staff. Complete policy and full list of medical necessity codes, refer to CMS website reference: <https://www.cms.gov/> ⇒

Flow Cytometry (CMS Article A56464)

CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367

polycythemia and basophilia are not sufficient to warrant FCM.

Lymphomas

In the current World Health Organization (WHO) classification, all non-Hodgkin lymphomas (NHLs) are distinct clinicopathologic entities defined by their clinical features, morphology, immunophenotype and, where appropriate, their genetic abnormalities. Immunophenotyping by FCM allows multiparameter evaluation of single cells and the ability to work on very small samples.

Most new cases of suspected NHL undergo initial immunophenotypic analysis as part of the routine handling of a specimen. A standard lymphoma panel is designed to identify abnormal populations of B cells, T cells and/or NK cells. A standard lymphoma panel might include a combination of markers from the following categories: T cells (CD2, CD3, CD4, CD5, CD7, CD8); B cells (CD19, CD20, CD23); Kappa and Lambda surface immunoglobulins light chains; plasma cells (CD38 and CD138); CALLA (CD10); CD45; CD56: FMC-7, CD103, CD11c, CD13, CD14, CD15, CD16 and CD34.

The immunophenotypes of lymphomas are widely known and FCM allows appropriate classification of most cases. However, atypical patterns occur and pose significant diagnostic difficulties where aberrant antigen expression patterns must be reconciled with morphology. Additional markers may be required to characterize the abnormal population of cells including markers of immature cells (HLA-DR), B cells (CD22) and myeloid cells (CD14, CD15, CD33, CD64, CD117).

Acute Leukemia

The diagnosis and management of acute leukemia depend on the detection, identification and characterization of leukemic cells. The identification of leukemic cells is straightforward in most occasions. However, each acute leukemia subgroup has heterogeneous biologic characteristics, many of which are associated with a different response to therapy.

As part of a routine diagnostic workup, most suspected acute leukemia cases undergo initial multiparameter immunophenotypic analysis, combined with morphology, cytochemistry, cytogenetics, and molecular biology.

A standard acute leukemia FCM panel is designed to determine whether leukemic blasts are of myeloid or lymphoid origin, and then to further classify the neoplastic cells (myeloid blasts, B lymphoblasts, abnormal promyelocytes, monoblasts, etc). An acute leukemia panel might include a combination of cell markers from the following categories: stem cell lineage (CD34), immature cell lineage (HLA-DR, CD 10); T cell (CD2, CD3, CD4, CD5, CD7 and CD8); B cell (CD19, CD20); myeloid cell (CD13, CD14, CD15, CD 33, CD64 and CD117); CD38, CD45, and CD56.

When the routine panel is insufficient to characterize the leukemic cells, additional antibodies including erythroid markers (CD71 and glycophorin A), megakaryocytic markers (CD41, CD61) or cytoplasmic markers may be indicated.

Chronic Lymphocytic Leukemia (CLL) & Other Chronic Lymphoproliferative Diseases (CLPD)

The history, physical exam (lymphadenopathy, splenomegaly and/or hepatomegaly) laboratory findings (lymphocytosis, granulocytopenia, anemia, thrombocytopenia), and lymphocyte morphology are suggestive of CLL. The diagnosis is established by paradoxical co-expression of CD5 on peripheral lymphocytes that express B cell markers (CD19, CD20, CD21 and CD 23) with Kappa or Lambda immunoglobulin light chain restriction. Additional markers such as CD38 and

REMINDER: The ordering provider is solely responsible for assigning diagnosis (codes). PDL does not, through this Reference Guide or otherwise, recommend any particular diagnosis codes. PDL will submit to Medicare only the diagnosis (codes) provided to PDL by the ordering provider and/or his/her authorized staff. Complete policy and full list of medical necessity codes, refer to CMS website reference: <https://www.cms.gov/> ⇒

Flow Cytometry (CMS Article A56464)

CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367

ZAP70 may provide important prognostic information.

FCM can distinguish CLL, the peripheral counterpart of small lymphocytic lymphoma, often diagnosed in lymph node biopsies, from other indolent lymphocytic malignancies including prolymphocytic leukemia, Waldenstrom's macroglobulinemia, leukemic phase of lymphomas, hairy cell leukemia, T-cell CLL, adult T-cell leukemia, large granulocytic leukemia and cutaneous T-cell lymphoma and natural killer (NK) disorders including KIR expression.

Plasma Cell Disorders

Plasma cell disorders are often identified through a combination of clinical, laboratory studies (urine or serum gamma globulins), morphologic, and radiologic findings. FCM immunophenotyping is useful to identify abnormal plasma cells, and the distinction between lymphoid and plasma cell neoplasms, and between reactive plasma cells and neoplastic plasma cells.

The initial FCM workup for a plasma cell disorder may include the basic lymphoma panel markers with additional markers such as CD28 and CD117.

Myelodysplastic Syndromes (MDS)

The gold standard for an MDS diagnosis is assessment of bone marrow smears for dysplastic changes. FCM may assist in MDS determination through the identification of abnormal maturing myeloid cells. An abnormal phenotype by FCM is a minimal diagnostic MDS criteria to establish a definitive diagnosis.

MDS has a definite risk and rate of progression to acute leukemia. Standard FCM leukemia panels are indicated to evaluate progression and onset of leukemia.

Chronic Myeloproliferative Disorders (CMPD)

Although genetic (Philadelphia chromosome and BCR/abl) and molecular studies (Jak 2) are the accepted cornerstone for the identification and classification of CMPDs, FCM may assist in the distinction from reactive hematopoietic proliferations and is important in the enumeration of blasts in the distinction from acute leukemia and an accelerated phase of CMPD.

CMPD also has a definite risk and rate of progression to acute leukemia. Standard FCM leukemia panels are indicated to evaluate progression and onset of leukemia.

Mast Cell Neoplasms

Mast cell neoplasms are uncommon disorders. Mast cells coexpress multiple markers including CD9, CD33, CD45, CD68, CD117, but also lack several myelomonocytic antigens including CD14, CD15, CD16 and most T- and B- cells antigens. Neoplastic mast cells have a similar antigen profile, but also can coexpress CD2 and CD25, which helps in distinguishing malignant mast cells from mastocytosis.

Paroxysmal hemoglobinuria (PNH)

PNH is a rare clonal hematopoietic disorder of stem cells. This condition is caused by genetic mutation that results in the

REMINDER: The ordering provider is solely responsible for assigning diagnosis (codes). PDL does not, through this Reference Guide or otherwise, recommend any particular diagnosis codes. PDL will submit to Medicare only the diagnosis (codes) provided to PDL by the ordering provider and/or his/her authorized staff. Complete policy and full list of medical necessity codes, refer to CMS website reference: <https://www.cms.gov/> ⇒

Flow Cytometry (CMS Article A56464)

CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367

absence of over a dozen surface antigens on red and white blood cells. FCM can diagnose PNH by assessing both the red and white blood cells for the absence of these antigens.

Minimal Residual Disease (MRD)

FCM analysis for MRD must identify phenotypic features characteristic of the disease of interest. The MRD flow analysis should not rely on an exact match between the phenotype of the residual disease and the original diagnostic specimen because phenotypes can change over time and with treatment. The antibody combinations should be chosen to maximize detection of disease, limit the impact of phenotypic variation, and permit detection of disease following antibody directed therapy.

Human Immunodeficiency Virus (HIV) Infection

HIV-1 infection causes significant changes in the number of CD4 and CD8 positive lymphocytes. CD4 count falls roughly 30% while CD8 count increases within 6 months after seroconversion, causing a decrease in the CD4/CD8 ratio.

Following HIV-1 diagnosis, FCM should include enumeration of mature T cells (CD3), helper T cells (CD4) and suppressor T cells (CD8) to ensure all major T cell subsets are accounted for (the sum of helper CD4 and suppressor CD8 T cells is roughly close to the total number of CD3 positive T cells). This ensures that the absolute CD4 is not artificially decreased due to sample degradation or other artifact.

A WBC count with differential also needs to be performed to calculate the absolute CD4 count (absolute lymphocyte count times CD4%).

Organ Transplants

In order to differentiate early rejection, immunosuppressive therapy toxicity or infection, FCM may be indicated to monitor postoperative organ transplants. CD3 is useful to monitor the effectiveness of certain immunosuppressive therapies. When the transplant patient demonstrates symptoms for the above conditions, repeated analysis may be required.

DNA Analysis

Carcinoma, Non-hematolymphoid Tumors

DNA analysis of tumor for ploidy and percent S-phase cells may be necessary for a few selective patients with carcinomas. When the obtained prognostic information will affect treatment decisions in patients with low stage (localized) disease, FCM results are useful.

Molar Pregnancy

FCM is useful to evaluate molar and partial molar pregnancies. Using a method to quantify DNA, similar to that used for evaluation of carcinomas, partial moles (triploid), can be distinguished from normal placenta and complete molar (diploid) pregnancies.

Primary Immunodeficiencies (PIDS)

PIDs are rare disorders that reflect inherited abnormalities in the development and maturation of cells responsible for

REMINDER: The ordering provider is solely responsible for assigning diagnosis (codes). PDL does not, through this Reference Guide or otherwise, recommend any particular diagnosis codes. PDL will submit to Medicare only the diagnosis (codes) provided to PDL by the ordering provider and/or his/her authorized staff. Complete policy and full list of medical necessity codes, refer to CMS website reference: <https://www.cms.gov/> ⇒

Flow Cytometry (CMS Article A56464)

CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367

immune function. More than 120 inherited immunodeficiency disorders are currently recognized. Affected individuals are prone to repeated infections, allergies, autoimmune disorders, and malignancies. Diagnosis typically occurs at an early age.

FCM may be indicated for diagnostic purposes and is usually limited to T (CD3, CD4, CD8), B (CD20) and NK cell (CD56) markers. Additional disease specific markers may be indicated.

Primary Platelet Disorders, Non-neoplastic

FCM is used for platelet analysis in quantitative and qualitative disorders such as Glanzmann Thrombasthenia (GT) and Bernard-Soulier Disease (B-S). GT is a rare inherited or acquired platelet disorder. Hereditary GT is defined by platelets with decreased expression or absence of the GPIIa/GPIIb receptor. This receptor is responsible for the initial platelet plug at the site of endothelial injury. Absence of the receptor may result in increased bleeding.

Acquired GT is likely an autoimmune phenomenon with the presence of GPIIb/GPIIIa blocking antibodies. FCM may be used to determine the functional effect and identify the molecular targets of these antibodies.

B-S is another rare inherited disorder that prevents the initial binding of platelets at the site of endothelial injury by absence of or presence of abnormal surface GPIa/V/IX receptor. Abnormalities of this receptor prevent attachment of platelets to subendothelial or free von Willebrand's factor with subsequent tendency to bleed.

FCM may be used to measure antibodies directed at specific loci of the GPIa/V/IX receptor, which include GPIb (CD42b), GPIX (CD42a), and GPV (CD42d). FCM is also used to assess the size of platelets in the initial evaluation of B-S disease. In B-S disease, platelets are generally larger than normal. FCM can distinguish B-S platelets from fragmented RBCs and debris by antibodies directed to the GPIb/IX/V receptor.

Red Cell and White Cell Disorders, Non-neoplastic

FCM is a valuable tool to establish abnormal or defective red blood cell, leukocyte and lymphocyte surface receptors, transmembrane molecules, and intracellular DNA.

It may be used in acquired and congenital red cell conditions such as in quantifying fetomaternal hemorrhage and hereditary spherocytosis, hereditary elliptocytosis, and hereditary persistence of fetal hemoglobin in the context of compound hemoglobinopathy syndromes.

FCM is a sensitive and specific method to identify leukocyte receptor abnormalities for the diagnosis of chronic granulomatous disease and CD11b deficiency.

It is an efficient method to identify lymphocytes HLA B27 associated with uveitis, ankylosing spondylitis, Reiter's syndrome and sacroiliitis.

Limitations:

Since FCM immunophenotypes for most common lymphomas and leukemias are well characterized, Noridian does **NOT** consider it "reasonable and necessary" to perform more than 24 markers in a panel. When atypical or unusual FCM results are obtained, the selective addition of more markers may be indicated.

REMINDER: The ordering provider is solely responsible for assigning diagnosis (codes). PDL does not, through this Reference Guide or otherwise, recommend any particular diagnosis codes. PDL will submit to Medicare only the diagnosis (codes) provided to PDL by the ordering provider and/or his/her authorized staff. Complete policy and full list of medical necessity codes, refer to CMS website reference: <https://www.cms.gov/> ⇒

Flow Cytometry (CMS Article A56464)

CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367

The flow report must document the specific indication for each marker over the 24-marker limit.

The FCM report must document the specific indication for each marker over the 24-marker limit. FCM reports without clear justification for each marker over 24 will be denied.

The ICD10 codes listed below are the top diagnosis codes currently utilized by ordering physicians for the limited coverage test highlighted above that are also listed as medically supportive under Medicare’s limited coverage policy.

If you are ordering this test for diagnostic reasons that are not covered under Medicare policy, an Advance Beneficiary Notice form is required.

ICD10 Codes commonly used for Flow Cytometry	
CPT Codes: 88182, 88184, 88185, 88187, 88188, 88189, 86355, 86356, 86357, 86359, 86360, 86361, 86367	
Code	Description
B20	Human immunodeficiency virus [HIV] disease
C32.1	Malignant neoplasm of supraglottis
C61	Malignant neoplasm of prostate
C73	Malignant neoplasm of thyroid gland
C76.0	Malignant neoplasm of head, face and neck
C79.51	Secondary malignant neoplasm of bone
C90.00	Multiple myeloma not having achieved remission
D64.9	Anemia, unspecified
D69.3	Immune thrombocytopenic purpura
D69.6	Thrombocytopenia, unspecified
D72.810	Lymphocytopenia
D72.818	Other decreased white blood cell count
D72.819	Decreased white blood cell count, unspecified
D72.829	Elevated white blood cell count, unspecified
D80.1	Nonfamilial hypogammaglobulinemia
D80.6	Antibody deficiency with near-normal immunoglobulins or with hyperimmunoglobulinemia
D83.1	Common variable immunodeficiency with predominant immunoregulatory T-cell disorders
D84.821	Immunodeficiency due to drugs
D89.89	Other specified disorders involving the immune mechanism, not elsewhere classified
Z21	Asymptomatic human immunodeficiency virus [HIV] infection status

REMINDER: The ordering provider is solely responsibility for assigning diagnosis (codes). PDL does not, through this Reference Guide or otherwise, recommend any particular diagnosis codes. PDL will submit to Medicare only the diagnosis (codes) provided to PDL by the ordering provider and/or his/her authorized staff. Complete policy and full list of medical necessity codes, refer to CMS website reference: <https://www.cms.gov/> ⇒